

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 318 144 A1**

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
11.06.2003 Bulletin 2003/24

(51) Int Cl.7: **C07C 403/16**

(21) Application number: **02023944.8**

(22) Date of filing: **25.10.2002**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
IE IT LI LU MC NL PT SE SK TR**
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• **Fehr, Charles**
1290 Versoix (CH)
• **Blanc, Pierre-Alain**
1263 Crassier (CH)

(30) Priority: **05.12.2001 WOPCT/IB01/02394**


(74) Representative:
Salvaterra-García, Maria de Lurdes
Firmenich SA,
P.O. Box 239
1211 Geneva 8 (CH)

(71) Applicant: **FIRMENICH SA**
CH-1211 Genève 8 (CH)

(54) Unsaturated ester as perfuming ingredient

(57) The present invention relates to the perfume industry. It concerns more particularly a compound of formula

wherein R represents a methyl or an ethyl group, in the form of any one of its isomers or of a mixture thereof. The invention relates also to the use of such a compound as a perfuming ingredient and to the perfumed article or perfuming composition containing a compound according to the invention.




(I)

EP 1 318 144 A1

Description

Technical field

[0001] The present invention relates to the perfume industry. It concerns more particularly a compound of formula



wherein R represents a methyl or an ethyl group, in the form of any one of its isomers or of a mixture thereof. The invention also relates to the use of such a compound as a perfuming ingredient and to the perfumed article or perfuming composition containing a compound according to the invention.


Prior art

[0002] Although several similar structures are known in the literature, to the best of our knowledge, the compounds of the present invention have novel chemical structures.

[0003] A 1/1 mixture of ethyl esters of formula (I) having the carbon-carbon double bond in position 3 or 4 of the ring is described in US 3,887,625. However in said document there are no useful indications that could allow a person skilled in the art to obtain an ester according to the invention (e.g. having the carbon-carbon double bond only in position 3). Furthermore, in the prior art there is no mention or suggestion of any organoleptic properties of the compounds of formula (I), or of any potential use of said compounds as perfuming ingredients.

Description of the invention

[0004] Surprisingly, we have now established that the compounds of formula



wherein R represents a methyl or an ethyl group, in the form of any one of its isomers, namely the (1R,2R), (1R,2S), (1S,2R) or the (1S,2S) isomers, or of a mixture thereof, possess a surprising and very useful damascone-like fragrance, which render them very convenient for the preparation of perfumes, perfuming compositions and perfumed products.

[0005] Amongst the compounds of formula (I), the methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate is much appreciated for its odor which presents a remarkable damascone, rosy note associated with citronellol like and slightly saffrony notes. The damascone note is particularly nice and has a very natural, very rosy, more precisely red rose odor. Moreover, said odor is devoid of the fermented-apple aspect present for example in the alpha damascone fragrance. From an olfactory point of view, the odor character of methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate is intermediate between that of delta and alpha damascone. Nevertheless, and despite this odor similarity, the use of the invention compounds has been proved to be more versatile than that of the damascenes.

[0006] Furthermore, perfumers have also noticed some differences between the two enantiomers of the trans methyl ester, namely the (1R,2S) and the (1S,2R) methyl 2,6,6-trimethyl-3-cyclohexene-1-carboxylate. Indeed, although the odor of both enantiomers is clearly damascone-like and in the same trend described hereinabove, the fragrance of the (1R,2S) enantiomer is stronger than the one of the (1S,2R) enantiomer which, in the other hand, is more aromatic.

[0007] In addition to the typical rose, damascone-like note of the compounds of formula (I), the methyl cis-2,6,6-trimethyl-3-cyclohexene-1-carboxylate possesses also terpenic and carrot tops notes.

[0008] The ethyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate develops a damascone-like odor similar to that of the corresponding trans methyl ester. The similarity of the odor of the ethyl and methyl ester is quite surprising in view of the prior art, wherein the methyl and ethyl esters of known analogues have in general significantly different odor properties.

[0009] The preferred compounds of the invention are the methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate and the (1R,2S) methyl 2,6,6-trimethyl-3-cyclohexene-1-carboxylate.

[0010] The character and the quality of the odor properties of the invention compounds are all the more surprising and unexpected in view of the odor of their prior art analogues.

[0011] Indeed, the trans methyl ester of formula (I) possesses an odor closer to that of damascones than its gamma or alpha isomers, namely methyl 2,2-dimethyl-6-methylene-1-cyclohexanecarboxylate (described in US 5,015,625, to Firmenich SA) and methyl 2,6,6-trimethyl-2-cyclohexene-1-carboxylate (described in EP 56109) respectively. This effect is mainly due to the fact that, when compared with the mentioned isomers, the trans methyl ester presents a more pure damascone-rosy connotation, which is almost devoid of the typical herbaceous-thujonic and red-fruity, green apple notes of said gamma and alpha isomer.

[0012] Similarly, when the trans ethyl ester of formula (I) is compared with the ethyl gamma isomer, also described in US 5,015,625, then the invention compound distinguishes itself from the prior known compound in that it presents a rosy odor devoid of the metallic, herbaceous-thujonic notes characteristic of the prior art analogue.

[0013] The compounds of the invention are suitable for use in fine perfumery, in perfumes, colognes or after-shave lotions, as well as in other current uses in perfumery such as to perfume soaps, preparations for the shower or the bath, such as bath salts, mousses, oils, gels or other preparations, products such as body oils, body-care products, body deodorants and antiperspirants, hair care products such as shampoos, ambient air deodorants, or cosmetic preparations.


[0014] The compounds of formula (I) can also be used in applications such as liquid or solid detergents for textile treatment, fabric softeners, or also in detergent compositions or cleaning products for cleaning dishes or varied surfaces, for industrial or household use.

[0015] In these applications, the compounds according to the invention can be used alone, as well as mixed with other perfuming ingredients, solvents, adjuvants or additives commonly used in perfumery. The nature and variety of these co-ingredients do not require a more detailed description here, which would not be exhaustive anyway. In fact, a person skilled in the art, having a general knowledge, is able to choose them according to the nature of the product that has to be perfumed and the olfactory effect sought. These perfuming co-ingredients belong to varied chemical groups such as alcohols, aldehydes, ketones, esters, ethers, acetates, nitrites, terpenic hydrocarbons, heterocyclic nitrogen- or sulfur-containing compounds, as well as natural or synthetic essential oils. Many of these ingredients are listed in reference texts such as S. Arctander, Perfume and Flavor Chemicals, 1969, Montclair, N.J., USA, or more recent versions thereof, or in other similar books, or yet in the specialized patent literature commonly available in the art.

[0016] The proportions in which the compounds according to the invention can be incorporated in the different products mentioned above vary in a broad range of values. Indeed, we have found that the compounds of the invention are environmentally friendly and not sensitizing, and this even at relatively high concentrations. The range of concentrations depends on the nature of the product to be perfumed and on the olfactory effect sought, as well as on the nature of the co-ingredients in a given composition when the compounds of the invention are used in admixture with perfuming co-ingredients, solvents or additives commonly used in the art.

[0017] For instance, concentrations from 0.1% to 1.5%, and preferably from 0.3% to 0.8%, by weight of these compounds, with respect to the perfuming composition in which they are incorporated, can be typically used. Lower concentrations than these can be used when these compounds are directly applied for perfuming some of the consumer products mentioned above.

[0018] The synthesis of the compounds of formula (I), which is another object of the present invention, is characterized by a rearrangement reaction which provides directly a final product having a carbon-carbon double bond selectively in the position 3. Said rearrangement reaction is characterized by the reaction of a α - β unsaturated tosyl hydrazone of formula



wherein R represents a methyl or ethyl group, in the presence of a borane derivative, such as NaBH₄ or catechol borane ((C₆H₄O₂)BH), and a carboxylate derivative selected in the group consisting of the C₁-C₈ carboxylic acids and their alkaline salts. Preferably the borane derivative is catechol borane and the carboxylate derivative is a sodium or potassium salt of a C₂-C₄ carboxylic acid.

[0019] This synthesis provides the compounds of formula (I) as a mixture of four possible isomers. The two diastereomers, namely the trans and cis isomers, may be subsequently separated by using conventional methods, such as recrystallization or chromatography. Finally, the enantiomers of each diastereomer may be separated by recrystallization of the ephedrate salt of the free acid derivative, as will be described in the examples.

[0020] The invention will now be described in further detail by way of the following examples, wherein the temperatures are indicated in degrees centigrade (°C); the NMR spectral data were recorded with a 360 MHz machine in CDCl₃; the chemical displacement δ are indicated in ppm with respect to the TMS as standard and all the abbreviations have the usual meaning in the art.

Example 1

[0021] Synthesis of methyl (trans/cis) 2,6,6-trimethyl-3-cyclohexene-1-carboxylate

a) Methyl 2,2,6-trimethyl-4-[(4-methylphenylsulfonyl)hydrazono]-2-cyclohexene-1-carboxylate

[0022] A suspension of p-toluenesulfonylhydrazide (96.80 g, 0.520 mol) in a solution of methyl 2,2,6-trimethyl-4-oxo-2-cyclohexene-1-carboxylate (92.71 g, 0.473 mol) and AcOH (0.463 g, 4.7 mmol) in MeOH (500 ml) was heated at reflux in a 1.5 l three necked flask, fitted with a mechanical stirrer.

[0023] Upon heating the reaction mixture became clear and, once at reflux temperature, the formed hydrazone gradually precipitated. Heating was prolonged for 6 h, then the reaction mixture was cooled at 0° and the precipitated solid was collected by filtration affording 153.80 g of pure hydrazone. Crystallization of the mother liquors from MeOH (55 ml) gave a second portion of pure hydrazone (2.90 g) (yield = 91%).

¹H-NMR: 0.92(s, 3H); 1.02(s, 3H); 1.80(s, 3H); 2.13(d, J=16.0, 1H); 2.39(d, J=16.0, 1H); 2.77(s, 3H); 3.66(s, 3H); 6.08(s, 1H); 7.29(d, J=8.0Hz, 2H); 7.84(d, J=8.0, 2H).
¹³C-NMR: 172.3(s); 154.1(s); 144.0(s); 140.3(s); 135.4(s); 129.6(d); 128.0(d); 124.5(d); 57.5(d); 51.92(q); 33.7(t); 33.4(s); 28.4(q); 27.3(q); 23.1(q); 21.6(q).

b) Methyl 2,6,6-trimethyl-3-cyclohexene-1-carboxylate and separation of trans and cis diastereomers

[0024] In a three necked 2 l flask fitted with a mechanical stirrer, catechol borane (15.25 g, 0.126 mol) was added in 15 minutes, at 0°C, to a suspension of the hydrazone obtained in a) (38.45 g, 0.105 mol) in CHCl₃ (500 ml). After 2 hours, to the yellow reaction mixture, still at 0°C, AcONa·3H₂O (28.7 g, 0.211 mol) was added at once. After 30 minutes, the reaction mixture was heated at reflux for 1 hour, then cooled at 25° and the white solid filtered off on Celite. The cake was washed with three portions of CHCl₃, and the combined filtrates were evaporated under reduced pressure. The crude material was dissolved in pentane, the organic layer washed (5% aqueous NaOH, H₂O and brine), dried (Na₂SO₄), filtered and concentrated. Bulb-to-bulb distillation (75-120°/3 mbar) of the crude product afforded 8.55 g of methyl 2,6,6-trimethyl-3-cyclohexene-1-carboxylate as a trans/cis = 86:14 mixture (yield = 43%).

[0025] Crystallization of the thus obtained product, at -78° from pentane (9.5 ml), afforded 6.39 g of methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate (trans/cis = 96:4), and concentration of the crystallization mother liquors gave 2.13 g of methyl 2,6,6-trimethyl-3-cyclohexene-1-carboxylate in the form of a trans/cis = 54:46 mixture. Two flash chromatograph separations (SiO₂, cyclohexane/AcOEt = 97:3) of the compounds issued from the mother liquors gave 0.9 g of the cis diastereomer as a trans/cis = 10:90 mixture.

Methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate

[0026]

¹H-NMR: 0.93(d, J=6.5, 3H); 0.95(s, 3H); 0.99(s, 3H); 1.74(d, J=18.0, 1H); 1.95(d, J=18.0, 1H); 2.04(d, J=11.5, 1H); 2.52(m, 1H); 3.69(s, 3H); 5.47(d, J=10.0, 1H); 5.52-5.59(m, 1H).
¹³C-NMR: 175.3(s); 131.1(d); 124.4(d); 57.6(d); 51.0(q); 41.1(t); 32.3(s); 31.6(d); 29.6(q); 20.9(q); 20.0(q).
 MS: 182(M⁺, 20); 151(10); 127(14); 126(16); 125(41); 123(66); 122(93); 114(13); 111(23); 108(17); 107(100); 96(13); 95(23); 93(22); 91(37); 83(19); 82(30); 81(43); 79(28); 77(27); 69(21); 68(28); 67(43); 65(16); 59(16); 55(21); 53(24); 51(10); 43(16); 39(41); 29(21); 27(22).

Methyl cis-2,6,6-trimethyl-3-cyclohexene-1-carboxylate

[0027]

¹H-NMR: 0.95(s, 3H); 0.97(s, 3H); 0.98(d, J=7.5, 3H); 1.66(d, J=18.0, 1H); 2.28(d, J=18.0, 1H); 2.37(d, J=6.0, 1H); 2.52(m, 1H); 3.61(s, 3H); 5.40(d, J=10.0, 1H); 5.65-5.73(m, 1H).
¹³C-NMR: 173.9(s); 128.2(d); 125.8(d); 54.6(d); 50.6(q); 35.6(t); 31.3(s); 29.6(d); 29.5(q); 29.0(q); 17.7(q).
 MS: 182(M⁺, 24); 167(10); 151(13); 127(16); 126(19); 125(43); 122(100); 114(14); 111(20); 108(15); 107(89); 95(18); 93(16); 91(25); 83(15); 82(28); 81(32); 79(16); 77(15); 69(18); 68(24); 67(30); 55(11); 53(10); 41(19); 39(11).

Example 2

[0028] Synthesis of the enantiomers of the methyl trans 2,6,6-trimethyl-3-cyclohexene-1-carboxylate

a) (±)-trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylic acid

[0029] A suspension of KOH (4.34 g, 77.5 mmol) in a solution of methyl trans 2,6,6-trimethyl-3-cyclohexene-1-carboxylate (13.48 g, 74.1 mmol) and thiophenol (8.51 g, 77.5 mmol) in DMF (60 ml) was heated at 100° in a three necked 250 ml flask fitted with a stirrer. After 16 hours, the cooled mixture was poured onto cold aqueous HCl (0.5 M, 200 ml) and extracted with Et₂O. The crude compound was pre-purified by standard acid-base extraction. The organic phase was dried (Na₂SO₄), filtered, concentrated, and then heated for 4 h at 100°/4 mbar to remove the excess of thiophenol. Bulb-to-bulb distillation (100-130°/0.1 mbar) afforded 10.57 g of the racemic acid as a trans/cis = 96:4 mixture (yield = 85%).

¹H-NMR: 1.02(s, 3H); 1.03(d, J=7.6Hz, 3H); 1.05(s, 3H); 1.77(d, J=18.0, 1H); 1.97(d, J=18.0, 1H); 2.04(d, J=10.0, 1s); 2.50(m, 1s); 5.47(d, J=10.0, 1H); 5.53-5.60(m, 1H); 11.35-11.85(s, 1H).
¹³C-NMR: 181.4(s); 130.9(d); 124.4(d); 57.6(d); 41.1(t); 32.2(s); 31.1(d); 29.5(q); 20.9(q); 20.0(q).
 MS: 168(M⁺, 91); 153(33); 125(34); 124(58); 113(58); 112(46); 111(100); 109(36); 108(20); 107(83); 101(13); 100(27); 97(26); 95(25); 91(36); 83(13); 82(65); 81(44); 79(24); 77(24); 69(58); 68(52); 67(60); 65(13); 56(18); 55(15); 53(13); 43(13); 41(24); 39(16).

b) (-)-trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylic acid

[0030] In a 100 ml flask equipped with a mechanical stirrer, a solution of (-)-ephedrine (Fluka, 5.64 g, 34.2 mmol) in an hexane-acetone mixture (75:25, 27 ml) was added in 5 minutes to a solution of (±)-trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylic acid (7.65 g, 45.5 mmol) in hexane (20 ml). After stirring for 45 minutes, the suspension was cooled at 0° and filtered to afford 5.61 g of ephedrate salt (65% ee) and a mother liquor. Two crystallizations of the ephedrate salt from a toluenehexane mixture (60:40) afforded 4.08 g of pure ephedrate salt and the corresponding mother liquors. The pure ephedrate salt thus obtained was dissolved in methanol (45 ml) and treated at 25° with 5% aqueous HCl (10.5 ml). After half an hour, the reaction mixture was concentrated and the residue dissolved in AcOEt (100 ml). The organic phase was washed (H₂O and brine), dried (Na₂SO₄), filtered and concentrated. A bulb-to-bulb distillation (100-130°/0.1 mbar) afforded 2.01 g of pure (-)-trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylic acid (98% ee determined by chiral GC on CP-Chirasil-DEX CB).

[α]_D²⁰ (CHCl₃, c = 14 g/l) = -97.3

NMR : as previously described for the racemic compound.

c) (+)-trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylic acid

[0031] The combined mother liquors obtained in b) (9.14 g of salt) were regenerated using the same procedure as described in b) to provide the free acid. A bulb-to-bulb distillation (100-130°C/0.1 mbar) afforded 5.20 g of (+)-trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylic acid (ee = 41%). The free acid was thus treated with (+)-ephedrine hemihydrate (Sigma, 5.39 g, 31.0 mmol) as described herein above for the other enantiomer to obtain 8.91 g of ephedrate salt (49% ee). Two crystallizations, as described in b), of the salt afforded 5.55 g of pure salt (97% ee). The pure ephedrate salt thus obtained was dissolved in methanol (58 ml) and treated with 5% aqueous HCl (14.5 ml). After half an hour, the reaction mixture was concentrated and the residue dissolved in AcOEt (100 ml). The organic phase was washed (H₂O and brine), dried (Na₂SO₄), filtered and concentrated. A bulb-to-bulb distillation (100-130°/0.1 mbar) afforded 2.01 g of pure (+)-trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylic acid (98% ee determined by chiral GC on

CP-Chirasil-DEX CB).

$[\alpha]_D^{20}$ (CHCl₃, c = 12 g/l) = +100.9

NMR : as previously described for the racemic compound.

5 d) (+) or (-)-methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate

10 [0032] In a three necked 100 ml flask, a solution of the desired enantiomer of free acid obtained in b) or c) (1.85 g, 11.0 mmol), methyl iodide (1.87 g, 15.8 mmol), and K₂CO₃ (1.82 g, 15.8 mmol) in DMF (40 ml) was stirred at 40°C for 2 hours. The cooled reaction mixture was poured onto cold 5% aqueous HCl and extracted twice with pentane. The organic phases were washed (three times with H₂O and brine), dried (Na₂SO₄), filtered and concentrated. A bulb-to-bulb distillation (100°/4 mbar) afforded 1.8 g of desired methyl ester (yield = 90%).

(-)-Methyl (1S,2R)-2,6,6-trimethyl-3-cyclohexene-1-carboxylate

15 [0033] $[\alpha]_D^{20}$ (CHCl₃, c = 28 g/l) = -83.3

NMR : as previously described for the racemic compound.

(+)-Methyl (1R,2S)-2,6,6-trimethyl-3-cyclohexene-1-carboxylate

20 [0034] $[\alpha]_D^{20}$ (CHCl₃, c = 29 g/l) = -83.1

NMR : as previously described for the racemic compound.

Example 3

25 Synthesis of ethyl trans 2,6,6-trimethyl-3-cyclohexene-1-carboxylate

30 [0035] In a three necked 100 ml flask a suspension of trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylic acid obtained in example 2 a) (0.800 g, 4.8 mmol), ethylbromide (0.616 g, 5.7 mmol) and K₂CO₃ (0.787 g, 5.7 mmol) in DMF (20 ml) was stirred at 40°C for 2 h. The cooled reaction mixture was next poured onto cold 5% aqueous HCl and extracted twice with pentane. The organic phases were washed (three times with H₂O and brine), dried (Na₂SO₄), filtered and concentrated. A bulb-to-bulb distillation (100°/4 mbar) afforded 0.870 g of ethyl trans 2,6,6-trimethyl-3-cyclohexene-1-carboxylate in the form of a trans/cis = 96:4 mixture (yield = 84%).

35 ¹H-NMR: 0.94(d, J=6.5, 3H); 0.96(s, 3H); 0.99(s, 3H); 1.28(t, J=7.6, 3H); 1.73(d, J=18.0, 1H); 1.96(d, J=18.0, 1H); 2.02(d, J=10.4, 1H); 2.52(m, 1H); 4.16(q, J=7.6, 2H); 5.46(d, J=10.0, 1H); 5.52-5.59(m, 1H).

¹³C-NMR : 174.7(s); 131.2(d); 124.3(d); 59.8(t); 57.7(d); 41.2(t); 32.3(s); 31.3(d); 29.4(q); 20.9(q); 19.9(q); 14.4(q).
MS: 196(23); 151(17); 139(18); 123(84); 122(100); 121(21); 112(25); 111(15); 108(14); 107(86); 95(15); 93(22); 91(30); 83(10); 82(22); 81(45); 79(23); 77(22); 69(15); 68(18); 67(32); 65(13); 55(18); 53(17); 42(18); 41(38); 39(26); 29(38); 27(24).

Example 4

45 Preparation of a perfuming composition

[0036] A "fruity" type base composition was prepared by admixing the following ingredients:

Ingredient	Parts by weight
10%* Amyl acetate	10
10%* 3-Methyl-2-butenyl acetate	30
10%* Aldehyde C6	30
Hexylcinnamic aldehyde	100
Ethyl anthranilate	5
2-Methyl-4-phenyl-2-butanol	30
Methyl cinnamate	10

* in dipropyleneglycol

(continued)

Ingredient	Parts by weight
10%* Cis-3-hexenol	30
Dihydro Eugenol	5
10%* β -Dorinone® 1)	20
Diethyl 1,4-cyclohexanedicarboxylate 2)	60
Habanolide® 3)	60
Hedione® 4)	120
Alpha ionone	50
10%* Cis-3-hexenol isobutyrate	20
Phenylethyl isobutyrate	30
10%* Cognac oil	20
Methyl isoeugenol	20
Muscenone Delta 5)	10
Floro® 6)	50
Phenethylol	150
Cis-3-hexenol salicylate	10
10%* Vanilline	25
Bergamot essential oil	40
Parnantheme 7)	5
	940

1) 1-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2-buten-1-one; origin: Firmenich SA, Geneva, Switzerland

2) origin : Firmenich SA, Geneva, Switzerland

3) pentadecenolide ; origin : Firmenich SA, Geneva, Switzerland

4) methyl dihydrojasmonate ; origin : Firmenich SA, Geneva, Switzerland

5) 3-methyl-(4 and 5)-cyclopentadecen-1-one; origin: Firmenich SA, Geneva, Switzerland

6) tetrahydro-2-isobutyl-4-methyl-4(2H)-pyranol; origin: Firmenich SA, Geneva, Switzerland

7) perfumery base ; origin : Firmenich SA, Geneva, Switzerland

[0037] The addition of 100 parts by weight of methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate to the above-described composition imparted to the latter a pronounced rose connotation. The olfactory effect provided by the addition of the invention compound is similar to that provided by δ -damascone. Moreover, this olfactory effect was less fruity-blackberry-blueberry than the one obtained by adding the same amount of the prior known gamma isomer (methyl 2,2-dimethyl-6-methylene-1-cyclohexanecarboxylate), and was also more elegant and more damascone-like than the effect that can be obtained with the other known structural analogues, e.g. the alpha isomer of the invention compound.

Example 5

Preparation of a perfuming composition

[0038] An "aromatic" type composition was prepared by admixing the following ingredients :

Ingredient	Parts by weight
Allyl amyl glycolate	30
Bergamot essential oil	100
Cetalox ® 1)	10
Coumarine	10
Dihydromyrcenol	300
Estragol	10
Exaltenone 2)	30

1) 8,12-epoxy-13,14,15,16-tetranorlabdane; origin: Firmenich SA, Geneva, Switzerland

2) (Z)-4-cyclopentadecen-1-one; origin : Firmenich SA, Geneva, Switzerland

EP 1 318 144 A1

(continued)

Ingredient	Parts by weight
10%* Galbanum essential oil	10
Hedione® HC ³⁾	70
Habanolide® ⁴⁾	80
10%* Isobutylquinoleine	40
Lavandin essential oil	40
Crystal moss	40
Polysantol® ⁵⁾	20
Patchouli essential oil	80
10%* Triplal ⁶⁾	40
Galbex® ⁷⁾ 183	60
	970

* in dipropylene glycol

3) methyl dihydrojasmonate ; origin : Firmenich SA, Geneva, Switzerland

4) pentadecenolide ; origin : Firmenich SA, Geneva, Switzerland

5) 3,3-dimethyl-5-(2',2',3'-trimethyl-3'-cyclopenten-1'-yl)-4-penten-2-ol ; origin : Firmenich SA, Geneva, Switzerland

6) origin : IFF, USA

7) origin : Firmenich SA, Geneva, Switzerland

[0039] The addition of 300 parts by weight of methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate to the above-described composition imparted to the latter a fruity, green, damascone-like, rosy odor. This effect was reminiscent of the fragrance generally imparted by delta damascone, but it was at the same time more fruity and less floral.

Example 6

Preparation of a perfuming composition

[0040] A "fruity-citrus" type base composition was prepared by admixing the following ingredients :

Ingredient	Parts by weight
Benzyl acetate	20
Geranyl acetate	20
Linalyl acetate	150
10%*2-Methyldecanal	20
Dihydromyrcenol	140
Habanolide® ¹⁾	20
Lemongrass	15
Lime	20
Limalol® ²⁾	5
Linalool	100
Hedione® ³⁾	60
Muscenone ⁴⁾	20
10%** (Z)-3-Decenal ⁵⁾	30

* in dipropylene glycol

** in ethyl citrate

1) pentadecenolide ; origin : Firmenich SA, Geneva, Switzerland

2) 1-p-menthene-9-carbaldehyde ; origin : Firmenich SA, Geneva, Switzerland

3) methyl dihydrojasmonate ; origin : Firmenich SA, Geneva, Switzerland

4) 3-methyl-(4 and 5)-cyclopentadecen-1-one ; origin : Firmenich SA, Geneva, Switzerland

5) origin : Firmenich SA, Geneva, Switzerland

(continued)


Ingredient	Parts by weight
10%* Oxane 6)	10
10%* Rose oxide	20
Orange essential oil	300
	950

6) cis-2-methyl-4-propyl-1,3-oxathiane; origin: Firmenich SA, Geneva, Switzerland

[0041] The addition of 50 parts by weight of methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate to the above-described base composition imparted to the latter a rosy and red apple effect. When the gamma or the alpha isomers were used instead of the compound of the invention, the effect was aromatic-rosemary and fruity-blueberry, or more cellar, herbaceous and earthy, respectively.

Claims


1. A compound of formula



(I)

wherein R represents a methyl or an ethyl group, in the form of any one of its isomers or of a mixture thereof.

2. As a compound according to claim 1, methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate, methyl cis-2,6,6-trimethyl-3-cyclohexene-1-carboxylate, ethyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate, methyl (1R,2S)-2,6,6-trimethyl-3-cyclohexene-1-carboxylate or methyl (1S,2R)-2,6,6-trimethyl-3-cyclohexene-1-carboxylate.
3. As a compound according to claim 1, methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate or methyl (1R,2S)-2,6,6-trimethyl-3-cyclohexene-1-carboxylate.
4. Use as a perfuming ingredient of a compound as defined in any one of claims 1 to 3.
5. A perfuming composition or a perfumed product comprising as perfuming ingredient a compound of formula



(I)

wherein R represents a methyl or an ethyl group, in the form of any one of its isomers or of a mixture thereof, together with a current perfuming co-ingredient, solvent or adjuvant.

6. A perfuming composition or a perfumed product according to claim 5, characterized in that the perfuming ingredient is methyl trans-2,6,6-trimethyl-3-cyclohexene-1-carboxylate or methyl (1R,2S)-2,6,6-trimethyl-3-cyclohexene-1-carboxylate.

EP 1 318 144 A1

7. A perfuming composition or a perfumed product according to claim 5 or 6, in the form of a perfume or a cologne, a perfumed soap, a shower or bath gel, a shampoo, a body deodorant or antiperspirant, an ambient air deodorant, a liquid or solid detergent for textile treatment, a detergent composition or a cleaning product for dishes or varied surfaces, a fabric softener or a cosmetic preparation.

5

10

15

20

25

30

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 02 02 3944

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	DE 32 33 175 A (FIRMENICH & CIE) 17 March 1983 (1983-03-17) * page 7 * * page 13 - page 14 *	1-3	C07C403/16
A	---	4-7	
X	DE 23 05 140 A (FIRMENICH & CIE) 16 August 1973 (1973-08-16) * page 9 * * page 13 * * example 4B *	1-3	
A	---	4-7	
X,D	& US 3 887 625 A (...) 3 June 1975 (1975-06-03) ---	1-3	
A	US 4 900 870 A (FEHR CHARLES ET AL) 13 February 1990 (1990-02-13) * abstract * -----	1-7	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C07C
Place of search		Date of completion of the search	Examiner
MUNICH		14 March 2003	Janus, S
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 (03.82) (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 02 3944

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

14-03-2003

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 3233175 A	17-03-1983	CH 648825 A5	15-04-1985
		DE 3233175 A1	17-03-1983
		JP 58065236 A	18-04-1983
		US 4460792 A	17-07-1984
DE 2305140 A	16-08-1973	CH 563951 A5	15-07-1975
		DE 2305140 A1	16-08-1973
		FR 2170238 A1	14-09-1973
		GB 1393719 A	14-05-1975
		GB 1393720 A	14-05-1975
		JP 48085557 A	13-11-1973
		NL 7301451 A ,B,	07-08-1973
US 4900870 A	13-02-1990	US 3887625 A	03-06-1975
		DE 3768592 D1	18-04-1991
		EP 0260472 A2	23-03-1988
		JP 2111485 C	21-11-1996
		JP 8019029 B	28-02-1996
		JP 63083039 A	13-04-1988

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82